



Outpatient Infusion Center  
Iron Order

Please fax form to: 580-585-5472

**Patient Information**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Gender: M F  
Patient Address: \_\_\_\_\_ Email: \_\_\_\_\_ Insurance: \_\_\_\_\_

**Additional Information Needed**

- Fax front/back of insurance card       Fax clinical/progress notes       Fax labs
- Fax patient demographics       Fax current medication list       Fax TB and Hep B results

**Diagnosis and Clinical Information**

**Diagnosis (ICD-10):**

- D50.0 Iron Deficiency Anemia Secondary to Blood Loss
- D50.8 Other Iron Deficiency Anemias
- D50.9 Iron Deficiency Anemia, Unspecified
- D63.1 Anemia in Chronic Kidney Disease
- Other: DX: \_\_\_\_\_

**Clinical Information:**

- New Therapy Induction       Therapy Change       Therapy Continuation
- Patient Weight: \_\_\_\_\_ lbs/ \_\_\_\_\_ kg       Patient Height: \_\_\_\_\_ in/ \_\_\_\_\_ cm
- Allergies: \_\_\_\_\_
- Therapies Tried and Failed: \_\_\_\_\_
- TB Test: Date: \_\_\_\_\_ Results: \_\_\_\_\_       Hep B Test: Date: \_\_\_\_\_ Results: \_\_\_\_\_
- Does the patient have non-dialysis dependent chronic kidney disease? Yes or No
- Is the patient currently on dialysis? Yes or No      Please indicate CKD Stage 1 2 3 4 5 Unknown
- Does the patient have venous access? Yes or No      If yes, What type? \_\_\_\_\_
- If No, initiate IV access.

**Lab Orders**

- CBC w/o diff     Ferritin     Iron/IBC
- Other: \_\_\_\_\_

**Lab Orders to be done by**

- Infusion Services
- Referring Provider

**Prescription Information**

- Injectafer       Dose: 15mg/kg (Patient weight <50kg)       Give 2 doses at least 7 days apart not to exceed 1500mg
- Dose: 750mg (Patient weight >50kg)       Give 2 doses at least 7 days apart not to exceed 1500mg
- Feraheme       Initial Dose: 510mg
- Second Dose: 510mg 3 to 8 days after initial dose
- Other       Medication: \_\_\_\_\_       Dose: \_\_\_\_\_       Frequency: \_\_\_\_\_

**Misc Orders**

- \_\_\_\_\_
- \_\_\_\_\_
- PICC/Midline/CAD dressing to be changed every 7 days

**Flushes**

- 10mL NS Flush Syringe PRN
- Heparin 500units/5mL Flush Syringe PRN
- 50ml NS Bag PRN
- 250ml NS Bag PRN

**Standing Orders for Adverse Reaction**

- Stop infusion and initiate NS bolus
- Notify Supervising physician and ordering provider
- Solu-Cortef 100mg SIVP signs of adverse reaction
- Benadryl 25 mg SIVP for hives or bronchial inflammation
- Epi 1:1000 1mL IM, IV or SQ for anaphylaxis
- Oxygen 2-5L nasal cannula
- Albuterol 2.5mg inhaled PRN to chest tightness
- Other: \_\_\_\_\_

**Prescriber Information**

Physician Name: \_\_\_\_\_ Official Contact Name: \_\_\_\_\_  
Contact #: \_\_\_\_\_ Fax Number: \_\_\_\_\_  
Address: \_\_\_\_\_ City/State/Zip: \_\_\_\_\_  
NPI#: \_\_\_\_\_ DEA#: \_\_\_\_\_ State License #: \_\_\_\_\_

Physician's Signature \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

