



Outpatient Infusion Center
Orencia Order

Please fax form to: 580-585-5472

Patient Information

Patient Name: DOB: Phone: Gender: M F
Patient Address: Email: Insurance:

Additional Information Needed

Fax front/back of insurance card Fax clinical/progress notes Fax labs
Fax patient demographics Fax current medication list Fax TB and Hep B results

Diagnosis and Clinical Information

Diagnosis (ICD-10):

- L40.52 Psoriatic Arthritis Multilans
L40.59 Other Psoriatic Arthropathy
M05.79 Rheumatoid Arthritis with Rheumatoid Factor of Multiple Sites without Organ or Systems Involvement
M05.89 Rheumatoid Arthritis with Rheumatoid Factor, multiple sites
M05.9 Rheumatoid Arthritis with Rheumatoid Factor, unspecified
M06.09 Rheumatoid Arthritis without Rheumatoid Factor, multiple sites
M06.89 Other Specified Rheumatoid Arthritis, multiple sets
M08.00 Unspecified Juvenile Rheumatoid Arthritis of Unspecified Site
M08.09 Unspecified Juvenile Rheumatoid Arthritis, Multiple sites

Other: DX: _____

Clinical Information:

New Therapy Induction Therapy Change Therapy Continuation
Patient Weight: lbs/ kg Patient Height: in/ cm
Allergies:
Therapies Tried and Failed:
TB Test: Date: Results: Hep B Test: Date: Results:
Does the patient have venous access? Yes or No If yes, What type?
If No, initiate IV access.

Lab Orders

CBC w/o diff CMP ESR CRP HBsAg HBsAB HbCAB Quantiferon Gold
Other: _____

Lab Orders to be done by

Infusion Services
Referring Provider

Prescription Information

Orencia Initial Dose: mg beginning week, week 2 and week 4
Maintenance Dose: mg every weeks after week 4
Weight <60kg= 500mg Weight 60-100kg=750mg Weight >100kg=1000mg

Misc Orders

PICC/Midline/CAD dressing to be changed every 7 days
Flashes
10mL NS Flush Syringe PRN
Heparin 500units/5mL Flush Syringe PRN
50ml NS Bag PRN
250ml NS Bag PRN

Standing Orders for Adverse Reaction

Stop infusion and initiate NS bolus Epi 1:1000 1mL IM, IV or SQ for anaphylaxis
Notify Supervising physician and ordering provider Oxygen 2-5L nasal cannula
Solu-Cortef 100mg SIVP signs of adverse reaction Albuterol 2.5mg inhaled PRN to chest tightness
Benadryl 25 mg SIVP for hives or bronchial inflammation Other: _____

Prescriber Information

Physician Name: Official Contact Name:
Contact #: Fax Number:
Address: City/State/Zip:
NPI#: DEA#: State License #:

Physician's Signature Date Time

