



Outpatient Infusion Center
Orencia Order

Please fax form to: 580-585-5472

Patient Information

Patient Name: _____ DOB: _____ Phone: _____ Gender: M F
Patient Address: _____ Email: _____ Insurance: _____

Additional Information Needed

- | | | |
|---|--|---|
| <input type="checkbox"/> Fax front/back of insurance card | <input type="checkbox"/> Fax clinical/progress notes | <input type="checkbox"/> Fax labs |
| <input type="checkbox"/> Fax patient demographics | <input type="checkbox"/> Fax current medication list | <input type="checkbox"/> Fax TB and Hep B results |

Diagnosis and Clinical Information

Diagnosis (ICD-10):

- L40.52 Psoriatic Arthritis Multilans
- L40.59 Other Psoriatic Arthropathy
- M05.79 Rheumatoid Arthritis with Rheumatoid Factor of Multiple Sites without Organ or Systems Involvement
- M05.89 Rheumatoid Arthritis with Rheumatoid Factor, multiple sites
- M05.9 Rheumatoid Arthritis with Rheumatoid Factor, unspecified
- M06.09 Rheumatoid Arthritis without Rheumatoid Factor, multiple sites
- M06.89 Other Specified Rheumatoid Arthritis, multiple sites
- M08.00 Unspecified Juvenile Rheumatoid Arthritis of Unspecified Site
- M08.09 Unspecified Juvenile Rheumatoid Arthritis, Multiple sites
- Other: DX: _____

Clinical Information:

- New Therapy Induction Therapy Change Therapy Continuation
- Patient Weight: _____ lbs/ _____ kg Patient Height: _____ in/ _____ cm
- Allergies: _____
- Therapies Tried and Failed: _____
- TB Test: Date: _____ Results: _____ Hep B Test: Date: _____ Results: _____
- Does the patient have venous access? Yes or No If yes, What type? _____
- If No, initiate IV access.

Lab Orders

- CBC w/o diff CMP ESR CRP HBsAg HBsAB HbcAB Quantiferon Gold
- Other: _____

Lab Orders to be done by

- Infusion Services
- Referring Provider

Prescription Information

- Orencia Initial Dose: _____ mg beginning week, week 2 and week 4
- Maintenance Dose: _____ mg every _____ weeks after week 4
- Weight <60kg= 500mg Weight 60-100kg=750mg Weight >100kg=1000mg

Misc Orders

- _____
- _____
- PICC/Midline/CAD dressing to be changed every 7 days
- Flushes**
- 10mL NS Flush Syringe PRN
- Heparin 500units/5mL Flush Syringe PRN
- 50ml NS Bag PRN
- 250ml NS Bag PRN

Standing Orders for Adverse Reaction

- Stop infusion and initiate NS bolus
- Notify Supervising physician and ordering provider
- Solu-Cortef 100mg SIVP signs of adverse reaction
- Benadryl 25 mg SIVP for hives or bronchial inflammation
- Epi 1:1000 1mL IM, IV or SQ for anaphylaxis
- Oxygen 2-5L nasal cannula
- Albuterol 2.5mg inhaled PRN to chest tightness
- Other: _____

Prescriber Information

Physician Name: _____ Official Contact Name: _____
 Contact #: _____ Fax Number: _____
 Address: _____ City/State/Zip: _____
 NPI#: _____ DEA#: _____ State License #: _____

Physician's Signature _____ Date _____ Time _____

