



Outpatient Infusion Center  
Osteoporosis Order

Please fax form to: 580-585-5472

**Patient Information**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Gender: M F

Patient Address: \_\_\_\_\_ Email: \_\_\_\_\_ Insurance: \_\_\_\_\_

**Additional Information Needed**

- ☐ Fax front/back of insurance card ☐ Fax clinical/progress notes ☐ Fax labs  
☐ Fax patient demographics ☐ Fax current medication list ☐ Fax TB and Hep B results

**Diagnosis and Clinical Information**

**Diagnosis (ICD-10):**

- ☐ M80.0 Age-Related Osteoporosis with current pathological fracture  
☐ M81.0 Age-related Osteoporosis without current pathological fracture  
☐ M81.8 Other Osteoporosis without current pathological fracture  
☐ M88.9 Paget's Disease of the bone in men and women  
☐ Other: DX: \_\_\_\_\_

**Clinical Information:**

- ☐ New Therapy Induction ☐ Therapy Change ☐ Therapy Continuation  
☐ Patient Weight: \_\_\_\_\_ lbs/ \_\_\_\_\_ kg ☐ Patient Height: \_\_\_\_\_ in/ \_\_\_\_\_ cm  
☐ Allergies: \_\_\_\_\_  
☐ Therapies Tried and Failed: \_\_\_\_\_  
☐ TB Test: Date: \_\_\_\_\_ Results: \_\_\_\_\_ ☐ Hep B Test: Date: \_\_\_\_\_ Results: \_\_\_\_\_  
☐ Is patient currently taking Calcium/ Vitamin D supplements? Yes or No Date of Last Calcium/ Vitamin D: \_\_\_\_\_  
☐ Does patient have a history of fractures? Yes or No  
☐ Date of last Dexa scan \_\_\_\_\_ ☐ Clinical Note for last DEXA scan attached? Yes or No  
☐ Does the patient have venous access? Yes or No If yes, What type? \_\_\_\_\_  
If No, initiate IV access.

**Lab Orders**

**Lab Orders to be done by**

- ☐ CBC w/o diff ☐ CMP ☐ ESR ☐ CRP ☐ HBsAg ☐ HBsAB ☐ HBcAB ☐ Quantiferon Gold ☐ Infusion Services  
☐ Other: \_\_\_\_\_ ☐ Referring Provider

**Prescription Information**

- ☐ Evenity ☐ Dose: 210mg (two 105mg injections) ☐ Frequency: Every month for 12 months  
☐ Prolia ☐ Dose: 60mg ☐ Frequency: Every 6 months  
☐ Zoledronic Acid ☐ Dose: 5mg ☐ Frequency: Once  
☐ Other ☐ Medication: \_\_\_\_\_ ☐ Dose: \_\_\_\_\_ ☐ Frequency: \_\_\_\_\_

**Misc Orders**

- ☐ \_\_\_\_\_ ☒ PICC/Midline/CAD dressing to be changed every 7 days

**Flushes**

- ☒ 10mL NS Flush Syringe PRN  
☒ Heparin 500units/5mL Flush Syringe PRN  
☒ 50ml NS Bag PRN  
☒ 250ml NS Bag PRN

**Standing Orders for Adverse Reaction**

- ☒ Stop infusion and initiate NS bolus ☒ Epi 1:1000 1mL IM, IV or SQ for anaphylaxis  
☒ Notify Supervising physician and ordering provider ☒ Oxygen 2-5L nasal cannula  
☒ Solu-Cortef 100mg SIVP signs of adverse reaction ☒ Albuterol 2.5mg inhaled PRN to chest tightness  
☒ Benadryl 25 mg SIVP for hives or bronchial inflammation ☐ Other: \_\_\_\_\_

**Prescriber Information**

Physician Name: \_\_\_\_\_ Official Contact Name: \_\_\_\_\_  
Contact #: \_\_\_\_\_ Fax Number: \_\_\_\_\_  
Address: \_\_\_\_\_ City/State/Zip: \_\_\_\_\_  
NPI#: \_\_\_\_\_ DEA#: \_\_\_\_\_ State License #: \_\_\_\_\_

Physician's Signature \_\_\_\_\_

Date \_\_\_\_\_

Time \_\_\_\_\_



PORD372

12/2011

Outpatient Infusion Center Osteoporosis Order

1 of 1