



Outpatient Infusion Center
Soliris Order

Please fax form to: 580-585-5472

Patient Information

Patient Name: DOB: Phone: Gender: M F
Patient Address: Email: Insurance:

Additional Information Needed

- Fax front/back of insurance card
Fax clinical/progress notes
Fax labs
Fax patient demographics
Fax current medication list
Fax TB and Hep B results

Diagnosis and Clinical Information

Diagnosis (ICD-10):

- D59.3 Atypical Hemolytic Uremic Syndrome (aHUS)
D59.5 Paroxysmal Nocturnal Hemoglobinuria (PNH)
G36.0 Neuromyelitis Optica Spectrum Disorder (NMOSD)
G70.00 Myasthenia Gravis (gMG) without Acute Exacerbation
G70.01 Myasthenia Gravis (gMG) with Acute Exacerbation
Other: DX:

Clinical Information:

- New Therapy Induction
Therapy Change
Therapy Continuation
Patient Weight: lbs/ kg
Patient Height: in/ cm
Allergies:
Therapies Tried and Failed:
TB Test: Date: Results:
Hep B Test: Date: Results:
Does the patient have a history of an adverse reaction to Soliris? Yes or No
Is the patient Anti-Acetylcholine Receptor (AChR) positive? (if gMG diagnosis) Yes or No
Is the patient Anti-Aquaporin Antibody positive? (if NMOSD diagnosis) Yes or No
Prescriber Enrolled in Soliris REMS Program? Yes or No
Patient Received Meningococcal Vaccine? Yes or No
Date of vaccination:
Does the patient have venous access? Yes or No
If yes, What type?

Lab Orders

- CBC w/o diff
CMP
ESR
CRP
Other:

Lab Orders to be done by

- Infusion Services
Referring Provider

Prescription Information

- Soliris
Initial Dose: 600mg weekly for first 4 weeks; then 900mg for 5th dose 1 week later
900mg weekly for the first 4 weeks; then 1200mg for 5th dose 1 week later
Maintenance Dose: 900mg every 2 weeks after initial dose
1200mg every 2 weeks after initial dose

Misc Orders

- PICC/Midline/CAD dressing to be changed every 7 days
Flushes: 10mL NS Flush Syringe PRN
Heparin 500units/5mL Flush Syringe PRN
50ml NS Bag PRN
250ml NS Bag PRN

Standing Orders for Adverse Reaction

- Stop infusion and initiate NS bolus
Notify Supervising physician and ordering provider
Solu-Cortef 100mg SIVP signs of adverse reaction
Benadryl 25 mg SIVP for hives or bronchial inflammation
Epi 1:1000 1mL IM, IV or SQ for anaphylaxis
Oxygen 2-5L nasal cannula
Albuterol 2.5mg inhaled PRN to chest tightness
Other:

Prescriber Information

Physician Name: Official Contact Name:
Contact #: Fax Number:
Address: City/State/Zip:
NPI#: DEA#: State License #:

Physician's Signature Date Time

